
Vitamin D and COVID-19

OPINIONS

GUNNAR HASLE

E-mail: hasle@reiseklinikken.com

Gunnar Hasle, specialist in infectious diseases, with a PhD on the intersection between medicine and zoology. Since 1999, he has been running 'Reiseklinikken' ('The Travel Clinic') in Oslo.

The author has completed the ICMJE form and declares no conflicts of interest.

To date, nobody has found a drug that dramatically improves the prospects in a severe case of SARS-CoV-2 infection, and it will be some time before a vaccine can become generally available. Could the vitamin D status of the population have a bearing on effective treatment and prevention of infections?

Vitamin D seems to play an important role for our immune system. A meta-analysis published in the British Medical Journal in 2017 concluded that vitamin D supplements gave general protection against respiratory infections, and the effect was most pronounced in those who had vitamin D deficiency [\(1\)](#). We see that many of the risk factors for COVID-19 (obesity, old age and dark skin pigmentation) coincide with the risk of vitamin D deficiency [\(2\)](#). In this context, it is notable that African-American patients have a higher risk of death from COVID-19 when compared to white patients [\(3\)](#), while COVID-19 does not appear to have affected Africa very severely [\(4\)](#). In Africa, the age distribution could be a partial explanation, but vitamin D from sunlight could be a contributory factor. The data are as yet sparse, but observational studies suggest that a low level of vitamin D increases the risk of catching COVID-19 [\(2\)](#). A recently published study compared available data on serum vitamin D in various countries with their incidence of and mortality from COVID-19 [\(5\)](#). The analysis indicated that higher levels of serum vitamin D made for a lower incidence of COVID-19, but there was insufficient evidence to draw conclusions

about an association between vitamin D levels and the degree of seriousness and mortality from COVID-19. Controlled trials are needed to establish whether vitamin D supplements have a prophylactic effect against COVID-19.

«Controlled trials are needed to establish whether vitamin D supplements have a prophylactic effect against COVID-19»

It is also conceivable that vitamin D supplements can be used in the treatment of COVID-19. A Spanish pilot study included hospitalised patients who were given chloroquine and azithromycin (which later has been proven to be ineffective against COVID-19) and randomised them for supplements of vitamin D capsules (266 µg). The treatment group (50 patients) received two capsules on the first day, one capsule on days 3, 5 and 7, and thereafter on a weekly basis (6). The 26 patients in the control group received no placebo. Only one of those 50 who received vitamin D needed intensive care and none of them died, whereas 13 in the control group needed intensive care and two died ($p < 0.001$). It will be interesting to see whether these findings can be replicated in randomised, placebo-controlled clinical trials. According to the authors, a large multi-centre study of vitamin D supplements is currently underway in Spain.

«In other words, irrespective of COVID-19, it is appropriate to recommend a daily spoonful of cod liver oil»

An obvious question is: Should everybody take cod liver oil? In 2014, a group of vitamin D experts appointed by the Nordic Council of Ministers recommended that the level should exceed 50 nmol/L (7). We have no reliable data on vitamin D levels in the Norwegian population, but a study from 2010 showed that the average value in non-smokers was a little higher than 50 nmol/L, and lower in the winter months (8). Vitamin D supplements are a simple, harmless and low-cost intervention, but will by no means render other measures against COVID-19 superfluous. The expert group recommended an intake of 10 µg/day, and 20 µg/day for the elderly. In other words, irrespective of COVID-19, it is appropriate to recommend a daily spoonful of cod liver oil (10 µg / 5 mL) or the equivalent, now that we are entering a season with less sunlight.

LITERATURE

1. Martineau AR, Jolliffe DA, Hooper RL et al. Vitamin D supplementation to prevent acute respiratory tract infections: systematic review and meta-analysis of individual participant data. *BMJ* 2017; 356: i6583. [PubMed] [CrossRef]
2. Martineau AR, Forouhi NG. Vitamin D for COVID-19: a case to answer? *Lancet Diabetes Endocrinol* 2020; 8: 735–6. [PubMed][CrossRef]
3. Price-Haywood EG, Burton J, Fort D et al. Hospitalization and mortality among black patients and white patients with Covid-19. *N Engl J Med* 2020;

382: 2534–43. [PubMed][CrossRef]

4. Worldometer. <https://www.worldometers.info/coronavirus/> Accessed 9.10.2020.

5. Ali N. Role of vitamin D in preventing of COVID-19 infection, progression and severity. *J Infect Public Health* 2020; 13: 1373–80. [PubMed][CrossRef]

6. Entrenas Castillo M, Entrenas Costa LM, Vaquero Barrios JM et al. "Effect of calcifediol treatment and best available therapy versus best available therapy on intensive care unit admission and mortality among patients hospitalized for COVID-19: A pilot randomized clinical study". *J Steroid Biochem Mol Biol* 2020; 203: 105751. [PubMed][CrossRef]

7. Brustad M, Meyer HE. Vitamin D–hvør mye er nok, og er mer bedre for helsen? *Tidsskr Nor Legeforen* 2014; 134: 726–8. [PubMed][CrossRef]

8. Grimnes G, Almaas B, Eggen AE et al. Effect of smoking on the serum levels of 25-hydroxyvitamin D depends on the assay employed. *Eur J Endocrinol* 2010; 163: 339–48. [PubMed][CrossRef]

Publisert: 9. November 2020. *Tidsskr Nor Legeforen*. DOI: 10.4045/tidsskr.20.0803

Received 7.10.2020, accepted 14.10.2020.

Copyright: © Tidsskriftet 2026 Downloaded from tidsskriftet.no 3 July 2026.