
Caesarean sections among immigrant women with different levels of education

ORIGINAL ARTICLE

HANNA SØLI OTTESEN

hannaottesen97@gmail.com

Institute of Health and Society

Faculty of Medicine

University of Oslo

Author contribution: idea, design, analysis, interpretation of data and preparation of the manuscript.

Hanna Søli Ottesen, 5th year medical student in the research programme for medical students

The author has completed the ICMJE form and declares no conflicts of interest.

INGVIL KRARUP SØRBYE

Norwegian Research Centre for Women's Health

Department of Obstetrics and Gynaecology

Oslo University Hospital, Rikshospitalet

and

Maternity ward

Oslo University Hospital, Rikshospitalet

Author contribution: idea, design, interpretation of data and revision of the manuscript.

Ingvil Krarup Sørbye, specialist in obstetrics and gynaecology, senior consultant.

The author has completed the ICMJE form and declares no conflicts of interest.

BENEDIKTE VICTORIA LINDSKOG

Department of Social Work, Child Welfare and Social Policy

Oslo Metropolitan University

Author contribution: interpretation of data and revision of the manuscript.

Benedikte Victoria Lindskog, associate professor.

The author has completed the ICMJE form and declares no conflicts of interest.

SIRI VANGEN

Norwegian Research Centre for Women's Health

Department of Obstetrics and Gynaecology

Oslo University Hospital, Rikshospitalet

and

Institute of Clinical Medicine

Faculty of Medicine

University of Oslo

Author contribution: interpretation of data and revision of the manuscript.

Siri Vangen, specialist in obstetrics and gynaecology, head of the Norwegian Research Centre for Women's Health and adjunct professor.

The author has completed the ICMJE form and declares no conflicts of interest.

JOHANNE SUNDBY

Institute of Health and Society

Faculty of Medicine

University of Oslo

Author contribution: design, interpretation of data and revision of the manuscript.

Johanne Sundby, specialist in obstetrics and gynaecology, professor emerita.

The author has completed the ICMJE form and declares no conflicts of interest.

KATRINE MARI OWE

Norwegian Research Centre for Women's Health

Department of Obstetrics and Gynaecology

Oslo University Hospital, Rikshospitalet

Author contribution: idea, design, analysis, interpretation of data and preparation and revision of the manuscript.

Katrine Mari Owe, researcher.

The author has completed the ICMJE form and declares no conflicts of interest.

BACKGROUND

Studies have shown a high incidence of emergency caesarean sections among immigrant women, especially those born in Sub-Saharan Africa, but the risk of planned and emergency caesarean section varies with the mother's level of education. The proportion of women with little or no education is higher among those born in Sub-Saharan Africa and other low- and middle-income countries than those born in Norway. We therefore wanted to investigate the relationship between maternal birthplace, level of education and risk of caesarean section.

MATERIAL AND METHOD

The study was based on all births recorded in the Medical Birth Registry of Norway between 2008 and 2017 linked to data from Statistics Norway. Maternal birthplace, divided into four categories, was the exposure variable. The outcome was planned or emergency caesarean section. We used multinomial logistic regression and stratified the analyses by level of education. Norwegian-born women constituted the reference group.

RESULTS

Of 572 349 births, immigrant women accounted for 26.6 %. Caesarean sections and emergency caesareans made up 15.1 % and 9.6 % of all births respectively. Norwegian-born women had the highest proportion of planned caesarean sections (5.7 %), while women born in Sub-Saharan Africa had the highest proportion of emergency caesareans (16.3 %). Among women with a higher education, the proportion of emergency caesareans was 8.3 % among Norwegian-born women and 18.1 % among women born in Sub-Saharan Africa (adjusted relative risk 2.41, 95 % confidence interval 2.18 to 2.66).

INTERPRETATION

The impact of education level on risk of caesarean section differed between immigrant women and Norwegian-born women.

Main findings

Women born in Sub-Saharan Africa and other low- and middle-income countries were found to have a high risk of emergency caesarean section regardless of reported education level compared to Norwegian-born women (adjusted relative risk from 1.06 to 2.41).

Among women with a higher education, those born in Sub-Saharan Africa had more than twice the risk of emergency caesarean section compared to Norwegian-born women (adjusted relative risk 2.41, 95 % CI 2.18 to 2.66).

Among women whose highest level of education was upper secondary school or below, those born abroad had a lower risk of planned caesarean section than Norwegian-born women (adjusted relative risk from 0.47 to 0.80).

According to the Norwegian Directorate of Health, the proportion of caesareans without a clear medical indication should be kept as low as possible (1). Caesarean section helps to reduce mortality and morbidity, but can lead to complications (2). The risk of complications is greater for emergency caesarean sections than for planned caesarean sections (3). In Norway, the proportion of caesarean sections has decreased slightly in the past ten years, from 16.5 % in 2011 to 15.8 % in 2020 (4).

Studies have shown that some immigrant groups have a significantly higher risk of caesarean section (5). This particularly applies to immigrant women born in Sub-Saharan Africa, whose caesarean section rate was 22.3 % between 2008 and 2018, compared to 15.6 % for Norwegian-born women (6, 7). This can only be partially explained by known medical risk factors such as obesity (8), gestational diabetes (9) and previous caesarean section. Non-medical factors such as education, health literacy, communication challenges and different cultural understandings of health and pregnancy also play an important role (10–12).

Previous studies have often excluded immigrant women (12, 13). We therefore wanted to investigate the relationship between maternal birthplace, education level and risk of emergency and planned caesarean section in Norway.

Material and method

Study population

We included all births in Norway between 2008 and 2017 (N = 608 980). A total of 36 631 births were excluded in accordance with the criteria shown in Figure 1. The study population consisted of 572 349 births, which were categorised according to maternal birthplace.




Figure 1 Exclusion criteria, study population and category of maternal birthplace.

Exposure

The exposure variable was maternal birthplace as recorded in the Norwegian Population Register. Maternal birthplace was categorised according to the WHO's Global Burden of Disease (GBD) framework, which consists of super-regions based on two criteria: geographic closeness and epidemiological similarity (14). Previous studies have shown that women born in Sub-Saharan Africa are a particularly vulnerable group, with a high incidence of adverse pregnancy outcomes (15), such as a high stillbirth rate (16) and a high risk of caesarean section (17). A large proportion of the women in this group also have little or no education (18, 19). Based on these findings, we wanted to investigate the relationship between education level and risk of caesarean section, with the main focus on women born in Sub-Saharan Africa. In order to draw comparisons with other immigrant populations, as well as women born in countries defined by the GBD index as high-income, we grouped the remaining GBD regions into the category '*other low- and middle-income countries*', referred to as '*low- and middle-income countries*' in the figures and tables. Norwegian-born women were used as the reference group (Figure 1).

Information on maternal level of education was based on the highest completed level of education, and was obtained from Norway's National Education Database, which contains education statistics at individual level dating back to 1970. Every ten years since 1991, Statistics Norway has collected information on completed education for immigrants in Norway for whom this information has not previously been reported (19).

The following four categories were used for education level: No completed education or below upper secondary education, Upper secondary education, Higher education and Unknown. Only 0.4 % had no completed education, and this group consisted solely of immigrant women. The proportion of immigrant women with unreported education was 18.4–21.3 % (Figure 2).




Figure 2 Distribution of maternal education level by maternal birthplace for 572 349 births between 2008 and 2017.

Outcome

Information on mode of delivery was obtained from the Medical Birth Registry of Norway and categorised as vaginal delivery, planned caesarean section or emergency caesarean section (defined as an emergency if the decision was made less than eight hours before the onset of labour). Unspecified caesareans ($n = 45$) were included in the emergency caesarean section group. The caesarean section variable in the Medical Birth Registry proved to have a high level of validity (20).

Other variables from the Medical Birth Registry

Maternal age and year of delivery were included as continuous variables, while body mass index (BMI) was categorised in line with the WHO classification. Health region was included due to differences in the caesarean section rate and immigrant population across hospital trusts. Women whose health region was not reported were placed in Southern and Eastern Norway Regional Health Authority. To reflect the differing risk of caesarean section between primiparous and multiparous women, parity was categorised as primiparous, multiparous without previous caesarean section or multiparous with previous caesarean section.

Statistical analyses

We used multinomial logistic regression analysis and present the results as relative risk (RR) with a 95 % confidence interval (CI). Norwegian-born women with a vaginal delivery formed the reference groups. The analyses were stratified by education level and adjusted for maternal age, year of delivery, health region, parity, and BMI. We wanted to examine the direct association between maternal birthplace and caesarean section in different educational strata and have therefore not adjusted for intermediate factors such as fetal position and pregnancy complications.

We carried out two sensitivity analyses to investigate the significance of fetal position and pregnancy complications for mode of delivery according to maternal birthplace. First, we only included births with normal cephalic presentation ($n = 522\ 116$). We then excluded pregnancy complications such as preeclampsia, gestational diabetes, chronic hypertension, and gestational hypertension, as these conditions increase the risk of caesarean section ($n = 524\ 612$).

All analyses were performed using STATA IC version 16 (Stata Statistical Software, College Station, TX, USA). The project was approved by the Regional Committees for Medical and Health Research Ethics (REK south-east, 2018/1086) and the Data Protection Officer at Oslo University Hospital (18 - 15786).

Results

The total study population included 572 349 births. Foreign-born women accounted for 26.6 % of the births: 4.8 % of the women were born in high-income countries, 3.8 % were born in Sub-Saharan Africa and 18.1 % were born in other low- and middle-income countries (Figure 1).

Table 1 describes the study population by mode of delivery. Caesarean sections accounted for 15.1 % of births, and 9.6 % were emergency caesarean sections. Norwegian-born women had the highest proportion of planned caesareans (5.7 %), while women born in Sub-Saharan Africa had the highest proportion of emergency caesareans (16.3 %). Women with unknown education had the highest proportion of emergency caesareans (11.3 %) and the lowest proportion of planned caesareans (4.7 %). A higher proportion of caesareans was seen for breech presentation, abnormal cephalic presentation, and pregnancy complications.

Table 1

Sociodemographic and obstetric variables for 572 349 births between 2008 and 2017 by mode of delivery. Number (%) if not otherwise specified. SD = standard deviation, BMI = body mass index.

Variable	Vaginal delivery (n = 485 875)	Emergency caesarean (n = 54 671)	Planned caesarean (n = 31 803)	Total (n = 572 349)
Maternal birthplace				
Norway	359 107 (85.5)	37 043 (8.8)	23 776 (5.7)	419 926 (73.4)
High-income country	22 972 (84.4)	2 777 (10.2)	1 483 (5.5)	27 232 (4.8)
Sub-Saharan Africa	17 090 (78.8)	3 534 (16.3)	1 076 (5.0)	21 700 (3.8)
Low- or middle-income country	86 706 (83.8)	11 317 (10.9)	5 468 (5.3)	103 491 (18.1)
Year of delivery				
2008–11	197 982 (84.7)	21 968 (9.4)	13 836 (5.9)	233 786 (40.9)
2012–14	145 090 (84.8)	16 545 (9.7)	9 421 (5.5)	171 056 (29.9)
2015–17	142 803 (85.3)	16 158 (9.7)	8 546 (5.1)	167 507 (29.3)
Age, mean (SD)	29.7 (5.1)	30.4 (5.3)	32.2 (5.1)	29.9 (5.2)
BMI category				
Underweight	12 575 (89.4)	926 (6.6)	573 (4.1)	14 074 (2.5)
Normal weight	182 523 (87.2)	17 118 (8.2)	9 721 (4.6)	209 362 (36.6)
Overweight	61 943 (82.7)	8 429 (11.3)	4 561 (6.1)	74 933 (13.1)
Obese	31 602 (77.5)	6 071 (14.9)	3 096 (7.6)	40 769 (7.1)
Unknown	197 232 (84.6)	22 127 (9.5)	13 852 (5.9)	233 211 (40.8)
Parity				
Primiparous	201 805 (82.9)	33 476 (13.8)	8 206 (3.4)	243 487 (42.5)
Multiparous	257 775 (93.8)	9 948 (3.6)	7 171 (2.6)	274 894 (48.0)
Multiparous with previous caesarean section	26 295 (48.7)	11 247 (20.8)	16 426 (30.4)	53 968 (9.4)
Education level				
Below upper secondary education	73 898 (83.8)	9 496 (10.8)	4 828 (5.5)	88 222 (15.4)
Upper secondary education	119 285 (84.1)	14 183 (10.0)	8 423 (5.9)	141 891 (24.8)

Variable	Vaginal delivery (n = 485 875)	Emergency caesarean (n = 54 671)	Planned caesarean (n = 31 803)	Total (n = 572 349)
Higher education	265 622 (85.7)	27 358 (8.8)	17 056 (5.5)	310 036 (54.2)
Unknown	27 070 (84.1)	3 634 (11.3)	1 496 (4.7)	32 200 (5.6)
Fetal position				
Normal cephalic presentation	459 731 (88.1)	39 082 (7.5)	23 303 (4.5)	522 116 (91.2)
Breech presentation	6 117 (30.4)	6 360 (31.6)	7 680 (38.1)	20 157 (3.5)
Abnormal cephalic presentation	19 804 (66.9)	9 019 (30.5)	795 (2.7)	29 618 (5.2)
Pregnancy complications				
Yes	33 936 (71.1)	10 319 (21.6)	3 482 (7.3)	47 737 (8.3)
No	451 939 (86.2)	44 352 (8.5)	28 321 (5.4)	524 612 (91.7)

Table 2 shows the unadjusted and adjusted relative risk for emergency and planned caesarean section by maternal birthplace, stratified by education level. Women with a higher education born in Sub-Saharan Africa had the highest risk of emergency caesarean section, with an adjusted relative risk of 2.41 (95 % CI 2.18 to 2.66).

Table 2

Unadjusted and adjusted relative risk (RR) with 95 % confidence interval (CI) for emergency and planned caesarean section by maternal birthplace, stratified by education level. Adjusted for health region, year of delivery, maternal age, parity and BMI. Norwegian-born women with a vaginal delivery form the reference group.

Education level	Maternal birthplace	Emergency caesarean			Planned caesarean		
		n (%)	RR (95%-CI)		n (%)	RR (95%-CI)	
			Unadjusted	Adjusted		Unadjusted	Adjusted
Below upper secondary education	Norway	5 078 (9.7)	Reference		3 052 (5.9)	Reference	
	High-income country	204 (10.5)	1.08 (0.93 to 1.25)	0.93 (0.79 to 1.08)	98 (5.1)	0.86 (0.70 to 1.06)	0.63 (0.51 to 0.79)
	Sub-Saharan Africa	1 700 (16.0)	1.74 (1.64 to 1.85)	1.59 (1.49 to 1.70)	476 (4.5)	0.81 (0.74 to 0.90)	0.47 (0.42 to 0.53)
	Low- or middle-income country	2 514 (10.7)	1.11 (1.05 to 1.16)	1.06 (1.01 to 1.12)	1 202 (5.1)	0.88 (0.82 to 0.94)	0.63 (0.58 to 0.68)
Upper secondary education	Norway	10 814 (9.5)	Reference		6 866 (6.0)	Reference	
	High-income country	537 (11.5)	1.24 (1.13 to 1.35)	1.08 (0.98 to 1.19)	261 (5.6)	0.95 (0.83 to 1.07)	0.77 (0.67 to 0.88)
	Sub-Saharan Africa	579 (17.4)	1.99 (1.82 to 2.19)	1.92 (1.74 to 2.12)	181 (5.4)	0.98 (0.84 to 1.14)	0.67 (0.57 to 0.79)
	Low- or middle-income country	2 253 (11.1)	1.18 (1.13 to 1.24)	1.17 (1.11 to 1.23)	1 115 (5.5)	0.92 (0.86 to 0.98)	0.80 (0.75 to 0.86)
Higher education	Norway	21 101 (8.3)	Reference		13 818 (5.5)	Reference	
	High-income country	1 506 (9.6)	1.18 (1.12 to 1.25)	1.04 (0.98 to 1.10)	885 (5.7)	1.06 (0.99 to 1.14)	0.88 (0.81 to 0.95)
	Sub-Saharan Africa	560 (18.1)	2.48 (2.26 to 2.72)	2.41 (2.18 to 2.66)	195 (6.3)	1.32 (1.14 to 1.53)	1.07 (0.91 to 1.26)
	Low- or middle-income country	4 191 (11.1)	1.38 (1.34 to 1.43)	1.31 (1.26 to 1.36)	2 158 (5.7)	1.09 (1.04 to 1.14)	1.06 (1.01 to 1.12)
Unknown education	Norway	50 (9.9)	Reference		40 (8.0)	Reference	
	High-income country	530 (10.6)	1.03 (0.76 to 1.40)	0.77 (0.56 to 1.06)	239 (4.8)	0.58 (0.41 to 0.82)	0.48 (0.32 to 0.71)
	Sub-Saharan Africa	695 (15.0)	1.55 (1.14 to 2.10)	1.58 (1.15 to 2.17)	224 (4.8)	0.62 (0.44 to 0.89)	0.48 (0.32 to 0.71)

Education level	Maternal birthplace	Emergency caesarean			Planned caesarean		
		n (%)	RR (95%-CI)		n (%)	RR (95%-CI)	
			Unadjusted	Adjusted		Unadjusted	Adjusted
	Low- or middle-income country	2 359 (10.7)	1.04 (0.77 to 1.40)	0.89 (0.66 to 1.21)	993 (4.5)	0.55 (0.39 to 0.76)	0.45 (0.31 to 0.66)

Among women with upper secondary education or below as their highest completed education, immigrant women had a lower risk of planned caesarean section than women born in Norway. Among those with a higher education, a slight difference in the risk of planned caesarean section was observed for women born in Sub-Saharan Africa and other low- and middle-income countries. The lowest risk of planned caesarean section was found in women with unknown education who were born in low- and middle-income countries (adjusted relative risk 0.45, 95 % CI 0.31 to 0.66).

Women who were born in Norway but whose education was unknown had a higher proportion of planned caesareans than other groups (8.0 %).

Figure 3 shows that among women with upper secondary education or below as their highest completed education, the risk of planned caesarean section was lower in foreign-born women than in Norwegian-born women. The risk of emergency caesarean section was significantly higher for women born in low- and middle-income countries, particularly women born in Sub-Saharan Africa, than for women born in Norway, regardless of education level.




Figure 3 Adjusted relative risk (RR) of emergency and planned caesarean section by maternal birthplace, stratified by maternal education level, for 572 349 births between 2008 and 2017. Adjusted for health region, year of delivery, maternal age, parity and BMI. Error bars represent 95 % confidence intervals. Norwegian-born women with a vaginal delivery constitute the reference group. Note the logarithmic scale.

Small changes were identified in the risk estimates when we limited the analyses to deliveries with normal cephalic presentation and when we excluded women with pregnancy complications (Tables 3 and 4).

Table 3

Deliveries with normal cephalic presentation ($n = 522\ 116$). Unadjusted and adjusted relative risk (RR) with 95 % confidence interval (CI) for emergency and planned caesarean section by maternal birthplace, stratified by education level. Adjusted for health region, year of delivery, maternal age, parity and BMI. Norwegian-born women with a vaginal delivery form the reference group.

Education level	Maternal birthplace	Emergency caesarean			Planned caesarean		
		n (%)	RR (95%-CI)		n (%)	RR (95%-CI)	
			Unadjusted	Adjusted		Unadjusted	Adjusted
Below upper secondary education	Norway	3 654 (7.7)	Reference		2 336 (4.9)	Reference	
	High-income country	135 (7.6)	0.99 (0.83 to 1.18)	0.85 (0.71 to 1.03)	73 (4.1)	0.84 (0.66 to 1.06)	0.62 (0.47 to 0.80)
	Sub-Saharan Africa	1 318 (13.5)	1.86 (1.74 to 1.99)	1.68 (1.56 to 1.81)	387 (4.0)	0.86 (0.77 to 0.95)	0.45 (0.40 to 0.51)
	Low- or middle-income country	1 850 (8.6)	1.13 (1.06 to 1.20)	1.08 (1.02 to 1.16)	889 (4.1)	0.85 (0.78 to 0.92)	0.57 (0.52 to 0.62)
Upper secondary education	Norway	7 715 (7.4)	Reference		5 172 (5.0)	Reference	
	High-income country	381 (9.0)	1.23 (1.11 to 1.37)	1.10 (0.98 to 1.23)	194 (4.6)	0.94 (0.81 to 1.08)	0.77 (0.65 to 0.90)
	Sub-Saharan Africa	457 (14.9)	2.18 (1.96 to 2.41)	2.13 (1.91 to 2.38)	139 (4.5)	0.99 (0.83 to 1.17)	0.65 (0.54 to 0.79)
	Low- or middle-income country	1 656 (9.0)	1.22 (1.15 to 1.29)	1.22 (1.15 to 1.29)	792 (4.3)	0.87 (0.80 to 0.94)	0.74 (0.68 to 0.80)
Higher education	Norway	14 778 (6.4)	Reference		9 905 (4.3)	Reference	
	High-income country	1 049 (7.4)	1.17 (1.10 to 1.25)	1.03 (0.96 to 1.10)	635 (4.5)	1.06 (0.98 to 1.15)	0.86 (0.79 to 0.95)
	Sub-Saharan Africa	436 (15.4)	2.71 (2.45 to 3.01)	2.67 (2.39 to 2.98)	159 (5.6)	1.48 (1.26 to 1.74)	1.20 (1.01 to 1.44)
	Low- or middle-income country	3 040 (8.8)	1.43 (1.37 to 1.48)	1.36 (1.31 to 1.42)	1 581 (4.6)	1.11 (1.05 to 1.17)	1.11 (1.04 to 1.18)

Education level	Maternal birthplace	Emergency caesarean			Planned caesarean		
		n (%)	RR (95%-CI)		n (%)	RR (95%-CI)	
			Unadjusted	Adjusted		Unadjusted	Adjusted
Unknown education	Norway	46 (9.8)	Reference		33 (7.1)	Reference	
	High-income country	358 (7.9)	0.75 (0.54 to 1.04)	0.53 (0.38 to 0.75)	151 (3.3)	0.44 (0.30 to 0.65)	0.33 (0.21 to 0.53)
	Sub-Saharan Africa	531 (12.4)	1.26 (0.92 to 1.74)	1.21 (0.87 to 1.69)	188 (4.4)	0.62 (0.42 to 0.92)	0.37 (0.23 to 0.59)
	Low- or middle-income country	1 678 (8.4)	0.80 (0.59 to 1.09)	0.64 (0.47 to 0.89)	669 (3.3)	0.44 (0.31 to 0.64)	0.31 (0.20 to 0.48)

Table 4

Births without pregnancy complications (preeclampsia, gestational diabetes, chronic hypertension and hypertension) ($n = 524\ 612$). Unadjusted and adjusted relative risk (RR) with 95 % confidence interval (CI) for emergency and planned caesarean section by maternal birthplace, stratified by education level. Adjusted for health region, year of delivery, maternal age, parity and BMI. Norwegian-born women with a vaginal delivery form the reference group.

Education level	Maternal birthplace	Emergency caesarean			Planned caesarean		
		n (%)	RR (95%-CI)		n (%)	RR (95%-CI)	
			Unadjusted	Adjusted		Unadjusted	Adjusted
Below upper secondary education	Norway	4 048 (8.5)	Reference		2 702 (5.7)	Reference	
	High-income country	162 (9.0)	1.05 (0.89 to 1.24)	0.89 (0.75 to 1.06)	90 (5.0)	0.88 (0.71 to 1.09)	0.63 (0.49 to 0.79)
	Sub-Saharan Africa	1 415 (14.7)	1.82 (1.71 to 1.95)	1.66 (1.55 to 1.78)	409 (4.2)	0.79 (0.71 to 0.88)	0.46 (0.41 to 0.52)
	Low- or middle-income country	2 024 (9.6)	1.12 (1.06 to 1.19)	1.08 (1.02 to 1.15)	1 008 (4.8)	0.84 (0.78 to 0.90)	0.61 (0.56 to 0.66)

Education level	Maternal birthplace	Emergency caesarean			Planned caesarean		
		n (%)	RR (95%-CI)		n (%)	RR (95%-CI)	
			Unadjusted	Adjusted		Unadjusted	Adjusted
Upper secondary education	Norway	8 563 (8.3)	Reference		6 068 (5.9)	Reference	
	High-income country	440 (10.4)	1.28 (1.15 to 1.41)	1.12 (1.01 to 1.24)	227 (5.3)	0.93 (0.81 to 1.06)	0.76 (0.66 to 0.88)
	Sub-Saharan Africa	468 (15.7)	2.06 (1.86 to 2.28)	1.96 (1.76 to 2.19)	154 (5.2)	0.96 (0.81 to 1.13)	0.65 (0.54 to 0.78)
	Low- or middle-income country	1 855 (10.1)	1.24 (1.17 to 1.30)	1.21 (1.15 to 1.28)	971 (5.3)	0.91 (0.85 to 0.98)	0.80 (0.74 to 0.86)
Higher education	Norway	17 112 (7.3)	Reference		12 506 (5.3)	Reference	
	High-income country	1 288 (8.9)	1.24 (1.17 to 1.31)	1.08 (1.02 to 1.15)	803 (5.5)	1.06 (0.98 to 1.14)	0.87 (0.80 to 0.94)
	Sub-Saharan Africa	447 (16.1)	2.45 (2.21 to 2.72)	2.41 (2.16 to 2.69)	157 (5.6)	1.18 (1.00 to 1.39)	0.99 (0.83 to 1.18)
	Low- or middle-income country	3 504 (10.2)	1.45 (1.39 to 1.50)	1.36 (1.31 to 1.41)	1 897 (5.5)	1.07 (1.02 to 1.13)	1.05 (0.99 to 1.11)
Unknown	Norge	42 (8.9)	Reference		36 (7.6)	Reference	
	High-income country	453 (9.7)	1.07 (0.77 to 1.50)	0.81 (0.58 to 1.15)	221 (4.8)	0.61 (0.42 to 0.88)	0.49 (0.33 to 0.75)
	Sub-Saharan Africa	572 (13.7)	1.57 (1.13 to 2.18)	1.61 (1.14 to 2.27)	178 (4.3)	0.57 (0.39 to 0.83)	0.43 (0.28 to 0.65)
	Low- or middle-income country	1 959 (9.7)	1.06 (0.77 to 1.46)	0.91 (0.65 to 1.27)	894 (4.4)	0.57 (0.40 to 0.80)	0.46 (0.31 to 0.68)

Discussion

The risk of emergency and planned caesarean section varied according to maternal birthplace and education level. Among Norwegian-born women, those with a higher education had the lowest incidence of both emergency and planned caesareans. Foreign-born women, particularly those born in Sub-Saharan Africa, had a high risk of emergency caesarean section regardless of education level. Among those with a lower level of education, foreign-born women had a lower risk of planned caesarean section compared to Norwegian-born women. This difference in risk estimates for planned caesarean section was not seen among women with a higher education.

Few studies have examined the relationship between maternal birthplace, education and the risk of caesarean section. A Norwegian study (12), which only included Norwegian-born women, found that the risk of both emergency and planned caesarean section was highest among women with the lowest level of education. This is consistent with our findings. However, we found higher risk estimates for emergency caesarean section among women from Sub-Saharan Africa and other low- and middle-income countries with a higher education, than for Norwegian-born women with the same level of education.

The incidence of planned caesarean sections was lower among foreign-born than Norwegian-born women but increased for foreign-born women in line with their education level. The difference in estimated risk for planned caesarean section between foreign-born and Norwegian-born women was less pronounced among those with a higher education. This has not been shown previously.

There may be several reasons why the education gradient had a differential impact on the risk of caesarean section among foreign-born women. Several studies have found a higher incidence of obesity (8), gestational diabetes (21) and previous caesarean sections among certain groups of immigrant women. Maternal BMI, parity and previous caesarean sections were all factored into our analyses. Adjustment for parity and previous caesarean sections affected the risk estimates, particularly for planned caesarean sections and for those with a low level of education. Adjusting for BMI led to small changes in our study, which is consistent with a recently published Norwegian study (22) showing a high risk of emergency caesarean section among women born in Sub-Saharan Africa in all BMI strata.

In addition to parity, maternal age had some significance for the risk estimates for both emergency and planned caesarean sections. When adjusting for maternal age, we accounted for the fact that maternal education level increases with age, regardless of maternal birthplace. Together with the results of the sensitivity analyses, this suggests that the high risk in women born in Sub-Saharan Africa cannot solely be explained by medical factors.

Studies have shown that non-medical factors such as language barriers, poorer health literacy and cultural attitudes to pregnancy can partly explain the higher risk of caesarean section in foreign-born women (7). Lack of communication

can adversely affect interaction during childbirth, and healthcare personnel may consequently fail to identify women with a medical indication for a planned caesarean section prior to delivery. The high proportion of emergency caesareans combined with the low proportion of planned caesareans may indicate underuse of planned caesareans among foreign-born women.

A long residence period can have a bearing on language skills and degree of integration. A recently published Norwegian study based on the same sample showed that women born in Sub-Saharan Africa with both long and short residence periods were at more than twice the risk of emergency caesarean section (22). Some immigrant women have poorer health literacy than women in the country they migrate to (23). This can impact on their confidence in the health service and treatment providers, use of healthcare services and understanding of symptoms of pregnancy complications. Different cultural understandings of pregnancy and modes of delivery among foreign-born women can also impact on the risk of caesarean section (24). Our study lacks information on various non-medical factors.

Strengths and limitations

The large study population is an important strength of this study. Using data from the Medical Birth Registry of Norway, where registration of all births is mandatory, enables a prospective approach with little likelihood of selection bias. Linking to other registries provides detailed information on maternal birthplace, education, and immigration background, and makes it possible to examine several potential confounding factors. It also enables analyses of subgroups without losing statistical power.

A significant weakness of the study was the high proportion of foreign-born women with unknown education (Figure 2). This group is included as a separate category in the analyses. Unknown education largely relates to education taken abroad, and this could lead to misclassification. Statistics Norway has carried out extensive work to obtain additional, correct information for this group (19). The group of Norwegian-born women with unknown education was small and had a higher proportion of both emergency and planned caesareans compared to other Norwegian-born women. No such difference was observed among foreign-born women, which means that the estimates for the group with unknown education should be interpreted with caution.

Unknown job status, occupation and income may have affected our results, as these variables are closely linked to socioeconomic status. Education is often used as a measure of socioeconomic status in Norway due to the country's comparatively small social differences and free health service. Nevertheless, education may not be an adequate measure of the socioeconomic status of immigrants (25, 26). This could be because the number of years of education does not necessarily correspond to a person's line of work or income, and this particularly applies to immigrants who end up in occupations with a low status and low pay as a result (18).

Conclusion

This study found that the risk of caesarean section varies according to maternal birthplace and education level, and that the impact of education level on risk of caesarean section differed between immigrant women and Norwegian-born women. Women born in low-income countries, particularly in Sub-Saharan Africa, have a high risk of emergency caesarean section regardless of education level, even when known risk factors are considered. The study sheds light on how maternal education as a measure of socioeconomic status affects pregnancy outcomes differently depending on where the mother was born.

This article has been peer-reviewed.

REFERENCES

1. Helsedirektoratet. Fødsel- keisersnitt 2021. <https://www.helsedirektoratet.no/statistikk/kvalitetsindikatorer/fodsel-og-abort/keisersnitt>. Accessed 3.2.2022.
2. Sandall J, Tribe RM, Avery L et al. Short-term and long-term effects of caesarean section on the health of women and children. *Lancet* 2018; 392: 1349–57. [PubMed][CrossRef]
3. Yang X-J, Sun S-S. Comparison of maternal and fetal complications in elective and emergency cesarean section: a systematic review and meta-analysis. *Arch Gynecol Obstet* 2017; 296: 503–12. [PubMed][CrossRef]
4. Folkehelseinstituttet. 2020-tallene fra Medisinsk fødselsregister 2020. <https://www.fhi.no/hn/helseregistre-og-registre/mfr/2020-tallene-fra-medisinsk-fodselsregister/> Accessed 30.11.2021.
5. Heslehurst N, Brown H, Pemu A et al. Perinatal health outcomes and care among asylum seekers and refugees: a systematic review of systematic reviews. *BMC Med* 2018; 16: 89. [PubMed][CrossRef]
6. Helsedirektoratet. Endring i fødepopulasjon og konsekvenser for bemanning og finansieringssystem. https://www.helsedirektoratet.no/rapporter/ending-i-fodepopulasjon-og-konsekvenser-for-bemanning-og-finansieringssystem/Rapport%20om%20f%C3%B8depopulasjonen.pdf/_/attachment/inline/3435df20-ea13-4d9f-99ed-f711d6ffbef0:51f3f1f4a94cd0893d94f09f3c7663d150ae61b0/Rapport%20om%20f%C3%B8depopulasjonen.pdf Accessed 30.1.2022
7. Merry L, Vangen S, Small R. Caesarean births among migrant women in high-income countries. *Best Pract Res Clin Obstet Gynaecol* 2016; 32: 88–99. [PubMed][CrossRef]

8. Gele AA, Mbalilaki AJ. Overweight and obesity among African immigrants in Oslo. *BMC Res Notes* 2013; 6: 119. [PubMed][CrossRef]
9. Gagnon AJ, McDermott S, Rigol-Chachamovich J et al. International migration and gestational diabetes mellitus: a systematic review of the literature and meta-analysis. *Paediatr Perinat Epidemiol* 2011; 25: 575–92. [PubMed][CrossRef]
10. Merry L, Semenic S, Gyorkos TW et al. International migration as a determinant of emergency caesarean. *Women Birth* 2016; 29: e89–98. [PubMed][CrossRef]
11. Esscher A, Binder-Finnema P, Bødker B et al. Suboptimal care and maternal mortality among foreign-born women in Sweden: maternal death audit with application of the 'migration three delays' model. *BMC Pregnancy Childbirth* 2014; 14: 141. [PubMed][CrossRef]
12. Tollånes MC, Thompson JM, Daltveit AK et al. Cesarean section and maternal education; secular trends in Norway, 1967-2004. *Acta Obstet Gynecol Scand* 2007; 86: 840–8. [PubMed][CrossRef]
13. Eriksen HS, Høy S, Irgens LM et al. Social inequalities in the provision of obstetric services in Norway 1967-2009: a population-based cohort study. *Eur J Public Health* 2020; 30: 491–8. [PubMed][CrossRef]
14. What is GBD and why is it important?
<https://www.healthdata.org/gbd/faq> Accessed 16.9.2022.
15. Bakken KS, Skjeldal OH, Stray-Pedersen B. Higher risk for adverse obstetric outcomes among immigrants of African and Asian descent: a comparison study at a low-risk maternity hospital in Norway. *Birth* 2015; 42: 132–40. [PubMed][CrossRef]
16. Vik ES, Aasheim V, Schytt E et al. Stillbirth in relation to maternal country of birth and other migration related factors: a population-based study in Norway. *BMC Pregnancy Childbirth* 2019; 19: 5. [PubMed][CrossRef]
17. Merry L, Small R, Blondel B et al. International migration and caesarean birth: a systematic review and meta-analysis. *BMC Pregnancy Childbirth* 2013; 13: 27. [PubMed][CrossRef]
18. Bye KS. Utdanning og lønnsnivå hos innvandrere.
<https://www.ssb.no/arbeid-og-lonn/artikler-og-publikasjoner/utdanning-og-lonnsniva-hos-innvandrere> Accessed 1.3.2022.
19. Statistisk sentralbyrå. Befolkningens utdanningsnivå 2022.
<https://www.ssb.no/utdanning/utdanningsniva/statistikk/befolkningens-utdanningsniva>. Accessed 20.8.2022.
20. Lehmann S, Baghestan E, Børdahl P et al. Validation of data in the Medical Birth Registry of Norway on delivery after a previous cesarean section. *Acta Obstet Gynecol Scand* 2017; 96: 892–7. [PubMed][CrossRef]

21. Gagnon AJ, Merry L, Haase K. Predictors of emergency cesarean delivery among international migrant women in Canada. *Int J Gynaecol Obstet* 2013; 121: 270–4. [PubMed][CrossRef]
 22. Jatta F, Sundby J, Vangen S et al. Association between Maternal Origin, Pre-Pregnancy Body Mass Index and Caesarean Section: A Nation-Wide Registry Study. *Int J Environ Res Public Health* 2021; 18: 5938. [PubMed][CrossRef]
 23. Villadsen SF, Hadi H, Ismail I et al. ehealth literacy and health literacy among immigrants and their descendants compared with women of Danish origin: a cross-sectional study using a multidimensional approach among pregnant women. *BMJ Open* 2020; 10: e037076. [PubMed][CrossRef]
 24. Essén B, Binder P, Johnsdotter S. An anthropological analysis of the perspectives of Somali women in the West and their obstetric care providers on caesarean birth. *J Psychosom Obstet Gynaecol* 2011; 32: 10–8. [PubMed][CrossRef]
 25. Racape J, Schoenborn C, Sow M et al. Are all immigrant mothers really at risk of low birth weight and perinatal mortality? The crucial role of socio-economic status. *BMC Pregnancy Childbirth* 2016; 16: 75. [PubMed][CrossRef]
 26. Blumenshine P, Egerter S, Barclay CJ et al. Socioeconomic disparities in adverse birth outcomes: a systematic review. *Am J Prev Med* 2010; 39: 263–72. [PubMed][CrossRef]
-

Publisert: 23. November 2022. Tidsskr Nor Legeforen. DOI: 10.4045/tidsskr.22.0256

Received 27.3.2022, first revision submitted 1.9.2022, accepted 10.10.2022.

Published under open access CC BY-ND. Downloaded from tidsskriftet.no 5 July 2026.