
Usefulness of measuring the serum concentration of antihypertensive drugs in uncontrolled hypertension

PERSPECTIVES

MIMI STOKKE OPDAL

uxmist@ous-hf.no

Mimi Stokke Opdal, MD, PhD, specialist in clinical pharmacology, associate professor at the Institute of Clinical Medicine, University of Oslo, and senior consultant at the Department of Pharmacology, Oslo University Hospital. The author has completed the ICMJE form and declares no conflicts of interest.

STINE ROGNSTAD

Stine Rognstad, MD, specialist in clinical pharmacology, senior consultant and PhD candidate at the Department of Pharmacology, Oslo University Hospital and the University of Oslo.

The author has completed the ICMJE form and declares no conflicts of interest.

FRODE EDVARDSEN

Frode Edvardsen, MD, specialist in general medicine and clinical pharmacology, general practitioner in Trondheim and senior consultant at the Department of Clinical Pharmacology, St Olav's Hospital, Trondheim University Hospital.

The author has completed the ICMJE form and declares no conflicts of interest.

LENE VERNÅS HALVORSEN

Lene Vernås Halvorsen, MD, senior consultant and PhD candidate at the Department of Nephrology, Oslo University Hospital and the University of Oslo.

The author has completed the ICMJE form and declares the following conflict of interest: she holds an unpaid position in the European Society of Hypertension.

RUNE MO

Rune Mo, MD, PhD, specialist in cardiology, senior consultant, section leader at the Clinic of Cardiology, St Olav's Hospital, Trondheim University Hospital, and associate professor at the Norwegian University of Science and Technology (NTNU).

The author has completed the ICMJE form and declares no conflicts of interest.

INGEBJØRG GUSTAVSEN

Ingebjørg Gustavsen, MD, PhD, specialist in clinical pharmacology, section leader and senior consultant at the Department of Pharmacology, Oslo University Hospital.

The author has completed the ICMJE form and declares no conflicts of interest.

ANNE CECILIE K. LARSTORP

Anne Cecilie K. Larstorp, MD, PhD, specialist in medical biochemistry, senior consultant at the Department of Medical Biochemistry, Oslo University Hospital, and associate professor at the Institute of Clinical Medicine, University of Oslo.

The author has completed the ICMJE form and declares no conflicts of interest.

CAMILLA LUND SØRAAS*

Camilla Lund Søråas, MD, PhD, specialty registrar in occupational medicine at the Section for Occupational and Environmental Medicine, Oslo University Hospital.

The author has completed the ICMJE form and declares no conflicts of interest.

ARNE HELLAND*

Arne Helland, MD, PhD, specialist in clinical pharmacology, senior consultant at the Department of Clinical Pharmacology, St Olav's Hospital, Trondheim University Hospital and associate professor at the Norwegian University of Science and Technology (NTNU).

The author has completed the ICMJE form and declares no conflicts of interest.

*Camilla Lund Søråas and Arne Helland have contributed equally to this article.

One-third of the adult population in Norway has hypertension. Despite receiving lifestyle advice and drug therapy, only half achieve the recommended blood pressure range. Measurements of serum drug concentrations can reveal poor medication adherence and can be used to personalise treatment.

The fact that half of patients with hypertension do not achieve their blood pressure goal despite receiving lifestyle advice and drug therapy is concerning (1–3). Persistent hypertension is associated with the development of diseases affecting the heart, arteries, brain and kidneys. Hypertension is the leading cause of years of life lost globally (4).

Various factors affect blood pressure, and monitoring techniques should therefore be standardised. In 2023, the European Society of Hypertension (ESH) published an updated version of the Guidelines for the Management of Arterial Hypertension, in which the definition of hypertension is based on the various methods of blood pressure monitoring, such as at a doctor's office, ambulatory (24-hour) and at home, where patients measure their own blood pressure (5), see Table 1.


Table 1

Definition of hypertension based on different ways of measuring average blood pressure. The Journal of Hypertension has consented to reproduction of this table (5).

Method	Systolic blood pressure (mmHg)		Diastolic blood pressure (mmHg)
Office BP, standardised monitoring	≥ 140	and/or	≥ 90
24-hour ambulatory BP			
24-hour period	≥ 130	and/or	≥ 80
Daytime	≥ 135	and/or	≥ 85
Nighttime	≥ 120	and/or	≥ 70
Home BP	≥ 135	and/or	≥ 85

The guidelines provide recommendations for treating hypertension with lifestyle advice and medication (5) (Figure 1). The roughly 50 % of patients who do not achieve their blood pressure goal (1–3) are defined as having uncontrolled hypertension. This may be due to various factors: improperly performed blood pressure monitoring, undiagnosed secondary hypertension, non-adherence to lifestyle advice, too few or insufficient doses of medications, unsuitable choice of medication, non-adherence to medication and/or ineffective pharmacotherapy. About 5 % of patients with hypertension have true resistant hypertension, defined as failure to control blood pressure despite taking maximum doses of three recommended medications, including a diuretic (5).

Drug therapy for hypertension in general practice



¹ If eGFR <45 ml/min/1.73 m², consider a loop diuretic instead of thiazide.

² For example loop diuretic, aldosterone antagonist, alpha blocker.

Figure 1 Recommended as first-line agents for the treatment of hypertension in the revised ESH guidelines. The figure has been slightly adapted and reproduced with the permission of the Journal of Hypertension (5). The figure is supplemented with our recommendations on the role of therapeutic drug monitoring in patient follow-up, in line with the recommendations in the guidelines. More detailed information on indications for assessing medication adherence can be found in the text of the article. ACEi = angiotensin-converting enzyme inhibitor, ARB = angiotensin II receptor blocker, CCB = dihydropyridine calcium channel blocker, BP = blood pressure.

Medications in blood pressure management

The guidelines recommend the use of two medications for most patients with hypertension, preferably as a two-drug single pill combination, to ensure adherence and achieve a synergistic effect on blood pressure (5), see Figure 1. Angiotensin-converting enzyme inhibitors (ACEi), angiotensin II receptor blockers (ARB), dihydropyridine calcium channel blockers (CCB) and thiazide diuretics are the recommended medications. Beta-blockers are also first-line treatment but are only recommended for use if there are additional indications, such as coronary disease, heart failure or tachyarrhythmias. In the case of resistant hypertension, a loop diuretic, potassium-sparing diuretic and/or alpha-blocker can be used in addition (5).

The various drug groups have well-documented effects on lowering blood pressure (6), morbidity and mortality. The health benefits of treatment are related to the reduction in blood pressure and less so to the specific medications used (5). However, a report from 2023 shows that the effect on systolic blood pressure of four different medications given to the same patient differed by up to 4.4 mmHg. This suggests that personalised drug therapy may improve treatment efficacy (7).

Pharmacological reasons for uncontrolled blood pressure

The lack of expected effect of medications on blood pressure can be due to pharmacokinetic and pharmacodynamic factors specific to the individual patient. Pharmacokinetic variation is caused by differences in the absorption, distribution, metabolism and excretion of a drug, which in turn will alter the concentration of the drug at the site of action, causing a changed drug response. Pharmacodynamic variation is caused by changes in the effect/mechanism of action of a drug without changing the serum concentration.

«The lack of expected effect of medications on blood pressure can be due to pharmacokinetic and pharmacodynamic factors specific to the individual patient»

Pharmacokinetic and pharmacodynamic variations can both be due to age, sex, medical history, altered organ function, drug interactions, environmental factors and pharmacogenetic factors specific to the patient. Lack of efficacy can also be due to a failure to adequately intensify or up-titrate treatment (referred to as 'physician inertia' in the ESH guidelines), or prescribing medications that are not suitable for the patient.

Are patients taking their medicine?

Various studies report that 10–30 % of patients with hypertension may have a low medication adherence (2, 8, 9). This could be due to adverse effects of their medication, a feeling that there is no point in taking the medicine, forgetfulness and poor routines, cognitive impairment or other illnesses that affect a patient's ability to follow a treatment regimen. For most patients, hypertension is asymptomatic, and this is associated with lower medication adherence compared to symptomatic conditions.

«For most patients, hypertension is asymptomatic, and this is associated with lower medication adherence compared to symptomatic conditions»

There are various ways to assess medication adherence: talking to the patient, directly observed therapy followed by blood pressure measurement, checking prescription refill rates and counting pills, using an electronic pill dispenser and measuring the concentration of the medication in serum or urine (10). The latter is considered the most objective and is also feasible in clinical practice.

Analyses to measure serum concentrations of antihypertensive drugs are available at St Olav's Hospital in Trondheim and at Oslo University Hospital (11, 12). The working group of the ESH recommends measuring serum or urine concentrations of antihypertensive drugs in patients who do not achieve their blood pressure goal (5, 13).

Measuring serum levels

The doctor can establish whether the patient is actually taking their medication and if the drug concentrations are in the expected range by measuring the serum level of antihypertensive drugs and comparing the results with the patient's medication list (14) (Table 2). The results can be assessed in conjunction with blood pressure readings and other clinical findings. The pharmacokinetic properties of antihypertensive drugs are well known. For example, altered kidney function, drug interactions and slow or fast metabolism due to changed pharmacogenetics can result in abnormal drug concentrations.

Table 2

Pharmacokinetic properties and dose-adjusted reference ranges for the most frequently prescribed antihypertensive drugs. Medications with low renal elimination are mainly excreted via bile/faeces. For four of the drugs, we measure the metabolite of the parent substance. A blood sample for determining serum concentration is taken before the morning dose at achieved steady-state concentration (after five half-lives). When using recommended doses, the result can be compared with the reference range in the far right column. A blood sample to check medication adherence can be taken at any time during the dosing interval, and a result below the lower limit of the reference range can be used as an indication of low adherence.

Antihypertensive drugs	Half-lives (t) ¹	Metabolism ²	Percentage of renal secretion ² (parent substance + metabolites)	Dose-related reference range (nmol/L) ²
Alpha-blockers				
Doxazosin (extended release)	15–19	CYP 3A4, (2D6, 2C9)	9	5–80
Beta-blockers				
Atenolol	6–9	None	50	75–750
Bisoprolol	10–12	CYP 2D6, (3A4)	100	10–200
Carvedilol	7–10	CYP 2D6, 2C9 etc.	< 2	2.5–50
Labetalol	5–8	Conjugation	65	50–1 000
Metoprolol (extended release)	3–4	CYP 2D6	95	10–500
Calcium channel blockers				
Amlodipine	35–50	CYP 3A4/5	70	10–40
Diltiazem (extended release)	4–10	CYP 3A4, (2D6)	70	100–500
Lercanidipine	8–10	CYP 3A4	50	0.20–5
Nifedipine (extended release)	6–11	CYP 3A4	80	20–150
Verapamil (extended release)	4–12	CYP 3A4 etc.	75	40–400
Angiotensin-converting enzyme inhibitors				
Enalaprilat (metabolite of enalapril)	11	Carboxylesterase	60	10–300
Lisinopril	12	None	100 unchanged	10–300
Ramiprilat (metabolite of ramipril)	13–17	Carboxylesterase	60	4–60
Angiotensin II receptor blockers				
Irbesartan	11–15	Conjugation	20	300–3 000
Candesartan	9	CYP 2C9 (low proportion)	30	15–200
Losartan carboxylic acid (metabolite of losartan)	6–9	CYP 2C9, 3A4 (low proportion)	35	30–350
Telmisartan	24	Conjugation	< 1	8–80

Antihypertensive drugs	Half-lives (t) ¹	Metabolism ²	Percentage of renal secretion ² (parent substance + metabolites)	Dose-related reference range (nmol/L) ²
Valsartan	6	CYP 2C9 (low proportion)	13 unchanged	300–4000
Thiazide diuretics				
Bendroflumethiazide	3	70 % (unknown enzymes)	30 unchanged	1.5–30
Hydrochlorthiazide	9–13	None	60–100 unchanged	15–300
Aldosterone antagonists				
Eplerenone	3–6	CYP 3A4	70	3.5–350
Canrenone (metabolite of spironolactone)	9–24	Unknown enzymes	60	15–300

¹Data from Thorstensen et al. (12)

²Data from Rognstad et al. (14)

The general idea with therapeutic drug monitoring is that if the serum concentration level is within the reference range, therapeutic effects can be expected. If the level is below this range, the dose can be increased, and if it is above, the dose can be reduced. Serum measurements can therefore serve as an aid for optimising the dosage and for choosing the most suitable medication for the patient. Measuring serum concentrations of antihypertensive drugs can help doctors identify and improve low medication adherence and adjust concentrations to dose-related reference ranges (2, 8, 9, 12, 15–17). However, more research is needed to determine whether such measurements have an effect on blood pressure, although some studies do suggest that they can help to control blood pressure (18).

Monitoring uncontrolled hypertension

Adherence to lifestyle advice and drug therapy are specifically addressed in the 2023 ESH guidelines (5). Assessment of medication adherence is recommended at all blood pressure checks, and particularly before changing medication, in secondary hypertension investigations and when true resistant hypertension is suspected. Adherence should also be assessed in patients with only a slight reduction in blood pressure despite combination therapy with two active ingredients. We recommend measuring drug concentrations to assess adherence in patients with uncontrolled hypertension (Figure 1) (19).

In order to ensure that analysis results present a representative picture of medication adherence, unannounced blood sampling is performed during routine check-ups. Blood sampling for testing can be carried out at any time within the dosing interval.

Information about the medication, dose, start/dose change date, date/time of last intake and blood sampling must be entered on the requisition form. To assess medication adherence over time, multiple blood samples must be taken, as most antihypertensive drugs have a relatively short half-life (Table 2).

«In order to ensure that analysis results present a representative picture of medication adherence, unannounced blood sampling is performed during routine check-ups»

In addition to identifying any non-adherence, serum concentration measurements can reveal pharmacokinetic abnormalities, which the doctor then takes into account in any potential dose adjustments or changes to medication (19). In such cases, the blood sampling for testing should be carried out before the morning dose is taken, and the test result should be compared with dose-related reference ranges (Table 2). Here too it is important to specify the medication, dose, start/dose change date, date/time of last intake and blood sampling on the requisition form.

Costs

The cost of cardiovascular medications, which are largely used to treat hypertension, was approximately NOK 1 billion in Norway in 2020 (20). Hypertension often requires lifelong medication, and inadequate treatment response also incurs high costs due to increased morbidity and the reduced capacity to work that is often entailed. Ensuring effective drug therapy is therefore important from both a medical and socioeconomic perspective (1). The Norwegian Health Economics Administration's rate for performing this type of therapeutic drug monitoring is currently around NOK 680. Patients typically use two medications and need at least one blood pressure check each year. The amount that a patient with uncontrolled hypertension pays for therapeutic drug monitoring is small compared to the actual cost. Few cost-benefit analyses have been conducted on antihypertensive drug monitoring, but one study concluded that there is a likely socioeconomic benefit (21).

Conclusion

Many patients with hypertension do not achieve their blood pressure goal, often due to low medication adherence. Serum concentration measurements of antihypertensive drugs can indicate whether a patient is taking their medication as prescribed and reveal pharmacokinetic abnormalities that might explain a reduced drug response. In line with the updated ESH guidelines, we recommend that this type of therapeutic drug monitoring is performed for patients with uncontrolled hypertension, including at blood pressure checks in first-line services.

The authors Opdal, Rognstad, Halvorsen, Mo, Gustavsen, Larstorp and Søråas have participated in the national hypertension study, IDA-studien.no.

REFERENCES

1. Global report on hypertension: the race against a silent killer.
<https://www.who.int/publications/i/item/9789240081062> Accessed 10.5.2024.
2. Bergland OU, Halvorsen LV, Søråas CL et al. Detection of Nonadherence to Antihypertensive Treatment by Measurements of Serum Drug Concentrations. *Hypertension* 2021; 78: 617–28. [PubMed][CrossRef]
3. Muntner P, Miles MA, Jaeger BC et al. Blood Pressure Control Among US Adults, 2009 to 2012 Through 2017 to 2020. *Hypertension* 2022; 79: 1971–80. [PubMed][CrossRef]
4. GBD 2019 Risk Factors Collaborators. Global burden of 87 risk factors in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet* 2020; 396: 1223–49. [PubMed][CrossRef]
5. Mancia G, Kreutz R, Brunström M et al. 2023 ESH Guidelines for the management of arterial hypertension The Task Force for the management of arterial hypertension of the European Society of Hypertension: Endorsed by the International Society of Hypertension (ISH) and the European Renal Association (ERA). *J Hypertens* 2023; 41: 1874–2071. [PubMed][CrossRef]
6. Wu J, Kraja AT, Oberman A et al. A summary of the effects of antihypertensive medications on measured blood pressure. *Am J Hypertens* 2005; 18: 935–42. [PubMed][CrossRef]
7. Sundström J, Lind L, Nowrouzi S et al. Heterogeneity in Blood Pressure Response to 4 Antihypertensive Drugs: A Randomized Clinical Trial. *JAMA* 2023; 329: 1160–9. [PubMed][CrossRef]
8. Berra E, Azizi M, Capron A et al. Evaluation of Adherence Should Become an Integral Part of Assessment of Patients With Apparently Treatment-Resistant Hypertension. *Hypertension* 2016; 68: 297–306. [PubMed][CrossRef]
9. Rabbia F, Fulcheri C, Di Monaco S et al. Adherence to antihypertensive therapy and therapeutic dosage of antihypertensive drugs. *High Blood Press Cardiovasc Prev* 2016; 23: 341–5. [PubMed][CrossRef]
10. Eskås PA, Heimark S, Eek Mariampillai J et al. Adherence to medication and drug monitoring in apparent treatment-resistant hypertension. *Blood Press* 2016; 25: 199–205. [PubMed][CrossRef]
11. Gundersen POM, Helland A, Spigset O et al. Quantification of 21 antihypertensive drugs in serum using UHPLC-MS/MS. *J Chromatogr B Analyt Technol Biomed Life Sci* 2018; 1089: 84–93. [PubMed][CrossRef]
12. Thorstensen CW, Clasen PE, Rognstad S et al. Development of UHPLC-MS/MS methods to quantify 25 antihypertensive drugs in serum in a cohort of patients treated for hypertension. *J Pharm Biomed Anal* 2022; 219. doi: 10.1016/j.jpba.2022.114908. [PubMed][CrossRef]

13. Endorsed by the European Society of Hypertension (ESH) Working Group on Cardiovascular Pharmacotherapy and Adherence. Nonadherence in Hypertension: How to Develop and Implement Chemical Adherence Testing. *Hypertension* 2022; 79: 12–23. [PubMed][CrossRef]
14. Rognstad S, Søråas CL, Bergland OU et al. Establishing Serum Reference Ranges for Antihypertensive Drugs. *Ther Drug Monit* 2021; 43: 116–25. [PubMed][CrossRef]
15. Halvorsen LV, Søråas CL, Larstorp ACK et al. Effect of Drug Monitoring on Adherence and Blood Pressure: A Multicenter Randomized Clinical Trial. *Am J Hypertens* 2024; 0. doi: 10.1093/ajh/hpae059. [PubMed][CrossRef]
16. Peeters LEJ, Feyz L, Boersma E et al. Clinical Applicability of Monitoring Antihypertensive Drug Levels in Blood. *Hypertension* 2020; 76: 80–6. [PubMed][CrossRef]
17. Peeters LEJ, Kappers MHW, Hesselink DA et al. Antihypertensive drug concentration measurement combined with personalized feedback in resistant hypertension: a randomized controlled trial. *J Hypertens* 2024; 42: 169–78. [PubMed][CrossRef]
18. Brinker S, Pandey A, Ayers C et al. Therapeutic drug monitoring facilitates blood pressure control in resistant hypertension. *J Am Coll Cardiol* 2014; 63: 834–5. [PubMed][CrossRef]
19. Edvardsen F, Øien CM, Helland A. Serumkonsentrasjonsmåling av blodtrykksmedisiner ved behandlingsresistent hypertensjon. *Tidsskr Nor Lægeforen* 2016; 136: 1514–5. [PubMed][CrossRef]
20. Reseptregisteret FHI. <https://www.fhi.no/he/reseptregisteret/> Accessed 10.5.2024.
21. Chung O, Vongpatanasin W, Bonaventura K et al. Potential cost-effectiveness of therapeutic drug monitoring in patients with resistant hypertension. *J Hypertens* 2014; 32: 2411–21, discussion 2421. [PubMed][CrossRef]

Publisert: 15. August 2024. *Tidsskr Nor Lægeforen*. DOI: 10.4045/tidsskr.24.0167

Received 21.3.2024, first revision submitted 20.4.2024, accepted 10.5.2024.

Copyright: © Tidsskriftet 2026 Downloaded from tidsskriftet.no 12 September 2026.